

7.5
1/6

Complex numbers.

$$z = a + bi$$

$|z|$ Modulus.

θ argument of z

$$z = r \cos \theta + i r \sin \theta$$

$$= r(\cos \theta + i \sin \theta). \quad (\text{polar form of } z)$$

Thm: De Moivre's Thm

$$(\cos \theta + i \sin \theta)^n = \cos(n\theta) + i \sin(n\theta)$$

Thm: F T o A

Any poly $p(\lambda)$ w/ complex coefficients splits, that is, it can be written as a product of linear factors.

$$p(\lambda) = k(\lambda - \lambda_1)(\lambda - \lambda_2) \dots (\lambda - \lambda_n)$$

for some complex numbers $\lambda_1, \dots, \lambda_n, \in k$.

De Moivre's Thm rotation & powers of z .

7.5

2/6

If $z = r(\cos\theta + i\sin\theta)$... then z^n spirals
in, out, or on the unit circle.

FToA Any poly $p(\lambda)$ w/complex coef. splits. That
is, it can be written as a product of
linear factors

$$p(\lambda) = k(\lambda - \lambda_1)(\lambda - \lambda_2) \cdots (\lambda - \lambda_n)$$

Cont. on next page.

7.5

3/6

ex1: Diagonalize $A = \begin{bmatrix} 11 & 6 \\ -15 & -7 \end{bmatrix}$

$$\det(A - \lambda I) = \begin{vmatrix} 11-\lambda & 6 \\ -15 & -7-\lambda \end{vmatrix}$$

$$= (11-\lambda)(-7-\lambda) + 90$$

$$= -77 - 4\lambda + \lambda^2 + 90$$

$$= \lambda^2 - 4\lambda + 13$$

Solve $0 = \lambda^2 - 4\lambda + 13$. $\lambda = \frac{4 \pm \sqrt{16 - 4(1)(13)}}{2(1)}$

$$= \frac{4 \pm \sqrt{-40}}{2}$$

$$= 2 \pm 3i$$

Find $\ker(\underbrace{A - (2+3i)I}_{})$

$$\begin{bmatrix} 9-3i & 6 \\ -15 & -9-3i \end{bmatrix} \xrightarrow[9-3i]{R_1 \rightarrow R_1}$$

$$\begin{bmatrix} 1 & \frac{3}{5} + \frac{1}{5}i \\ -15 & -9-3i \end{bmatrix} \xrightarrow{15R_1 + R_2 \rightarrow R_2}$$

$$\begin{bmatrix} 1 & \frac{3}{5} + \frac{1}{5}i \\ 0 & 0 \end{bmatrix}$$

$$E_{2+3i} = \text{span}\left(\begin{bmatrix} \vec{v}_1 \\ -3-i \\ 5 \end{bmatrix}\right)$$

and check if the conjugate of \vec{v}_1 is also an eigenvector. $\lambda_2 \vec{v}_2$

$$\begin{bmatrix} 11 & 6 \\ -15 & -7 \end{bmatrix} \begin{bmatrix} -3+i \\ 5 \end{bmatrix} = \begin{bmatrix} -3+11i \\ 10-15i \end{bmatrix}$$

7.5
4/6

So $\begin{bmatrix} 2+3i & 0 \\ 0 & 2-3i \end{bmatrix} = P^{-1} A P$ (of the form $D = S^{-1}AS$)

where $P = \begin{bmatrix} -2-i & -3+i \\ 5 & 5 \end{bmatrix}$

cont. on next page.

ex 2: Diagonalize $B = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ (rotation-scaling matrix)

$$\lambda = a \pm ib$$

$a, b \in \mathbb{R}$ and $b \neq 0$.

$$E_{a+ib} = \text{span} \begin{bmatrix} i \\ 1 \end{bmatrix}$$

$$E_{a-ib} = \text{span} \begin{bmatrix} -i \\ 1 \end{bmatrix}$$

and $R^{-1} \begin{bmatrix} a & -b \\ b & a \end{bmatrix} R = \begin{bmatrix} a+ib & 0 \\ 0 & a-ib \end{bmatrix}$

(of the form $S^{-1}AS = D$).

recall: IF A is a real 2×2 matrix w/ eigenvalues $a \pm ib$ (b $\neq 0$) and corresponding eigenvectors $\vec{v} \pm \vec{w}$, then

$P^{-1} A P = \begin{bmatrix} a+ib & 0 \\ 0 & a-ib \end{bmatrix}$ where $P = \begin{bmatrix} 1 & 1 \\ \vec{v} + \vec{w} & \vec{v} - \vec{w} \end{bmatrix}$

matrix w/
complex
eigenvals.
 \downarrow

$$\Rightarrow P^{-1} A P = R^{-1} \begin{bmatrix} a & -b \\ b & a \end{bmatrix} R$$

$$\Rightarrow \begin{bmatrix} a & -b \\ b & a \end{bmatrix} = R P^{-1} A P R^{-1} = S^{-1} A S$$

where $S = P R^{-1} = \begin{bmatrix} c & d \\ e & f \end{bmatrix}$

S has real values \Leftrightarrow S is similar to a rotation-scaling matrix.

7.5
6/6

ex1 rev: $A = \begin{bmatrix} 11 & 6 \\ -15 & -7 \end{bmatrix}$

$$\lambda = 2 \pm 3i$$

$$E_{2+3i} = \text{span} \left[\underbrace{\begin{bmatrix} -3 & -i \\ 5 & \end{bmatrix}}_{\begin{bmatrix} -3 \\ 5 \end{bmatrix} + i \begin{bmatrix} -1 \\ 0 \end{bmatrix}} \right]$$

$$\vec{v} + i\vec{w}$$

$$\Rightarrow S = \begin{bmatrix} -1 & -3 \\ 0 & 5 \end{bmatrix}$$

and $S^{-1} A S = \begin{bmatrix} 2 & -3 \\ 3 & 2 \end{bmatrix} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$

where $\lambda = a+ib$ is an eigenvalue.

what scaling factor?

what notation?

Thm: A complex $n \times n$ matrix has n complex eigenvalues if they are counted w/ alg. mult.

Thm: $\det(A) = \lambda_1 \cdots \lambda_n$

$\text{Tr}(A) = \lambda_1 + \dots + \lambda_n$,

CHANGE OF BASIS MATRICES REQUIRE COMPLEX CONJUGATE EIGENVECTORS

$\lambda = 2 \pm 3i$ for the matrices $A = \begin{bmatrix} 11 & 6 \\ -15 & -7 \end{bmatrix}$ and $P = \begin{bmatrix} -3-i & -3+i \\ 5 & 5 \end{bmatrix}$.
 Eigenvectors of $B = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ can take many forms.

$$\vec{v}_1 = \begin{bmatrix} i \\ 1 \end{bmatrix}$$

$$\vec{v}_1 \text{ conjugate} \quad *(-1)$$

$$\vec{v}_2 = \begin{bmatrix} -i \\ 1 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} i \\ -1 \end{bmatrix}$$

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$

$$*(i)$$

$$\begin{bmatrix} 1 \\ i \end{bmatrix}$$

$$*(2)$$

$$\begin{bmatrix} -2i \\ 2 \end{bmatrix}$$

$$\begin{bmatrix} 1 \\ i \end{bmatrix}$$

or

which gives R many forms

$$R = \begin{bmatrix} i & -i \\ 1 & 1 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} i & i \\ 1 & -1 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} i & 1 \\ 1 & i \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} i & -2i \\ 1 & 2 \end{bmatrix}$$

check R & P by comparing the product

$$S = PR^{-1} \text{ to the formula for } S = \begin{bmatrix} \vec{v} & \vec{w} \end{bmatrix}$$

$$\text{where the eigenvectors of } A = \vec{v} + i\vec{w} = \begin{bmatrix} -3 \\ 5 \end{bmatrix} + i \begin{bmatrix} 4 \\ 0 \end{bmatrix}$$

$$\vec{v}_1 \text{ conjugate: } PR^{-1} = \frac{1}{2i} \begin{bmatrix} -3-i & -3+i \\ 5 & 5 \end{bmatrix} \begin{bmatrix} 1 & i \\ -1 & i \end{bmatrix} = \frac{1}{2i} \begin{bmatrix} -2i & -6i \\ 0 & 10i \end{bmatrix} = \begin{bmatrix} -1 & -3 \\ 0 & 5 \end{bmatrix} = S$$

$$*(-1): PR^{-1} = -\frac{1}{2i} \begin{bmatrix} -3-i & -3+i \\ 5 & 5 \end{bmatrix} \begin{bmatrix} -1 & -i \\ -1 & i \end{bmatrix} = +\frac{1}{2i} \begin{bmatrix} -6 & +2 \\ +10 & 0 \end{bmatrix} = \begin{bmatrix} 3i & -i \\ -5i & 0 \end{bmatrix} \neq S$$

$$*(i): PR^{-1} = -\frac{1}{2} \begin{bmatrix} -3-i & -3+i \\ 5 & 5 \end{bmatrix} \begin{bmatrix} i & -1 \\ -1 & i \end{bmatrix} = -\frac{1}{2} \begin{bmatrix} -4i+4 & 2-2i \\ 5i-5 & -5+5i \end{bmatrix} \neq S$$

$$*(2): PR^{-1} = \frac{1}{4i} \begin{bmatrix} -3-i & -3+i \\ 5 & 5 \end{bmatrix} \begin{bmatrix} 2 & 2i \\ -1 & i \end{bmatrix} = \frac{1}{4i} \begin{bmatrix} -3-2i & -4i+1 \\ 5 & 15i \end{bmatrix} \neq S$$

THE FORMULA FOR S

$$\text{If } R = \begin{bmatrix} i & -i \\ 1 & 1 \end{bmatrix} \text{ and } P = \begin{bmatrix} 1 & 1 \\ \bar{v} + i\bar{\omega} & v - i\bar{\omega} \end{bmatrix}$$

$$\begin{aligned} \text{Then } S &= PR^{-1} = \frac{1}{2i} \begin{bmatrix} 1 & 1 \\ \bar{v} + i\bar{\omega} & v - i\bar{\omega} \end{bmatrix} \begin{bmatrix} 1 & i \\ -1 & i \end{bmatrix} \\ &= \frac{1}{2i} \begin{bmatrix} a+ib & a-ib \\ c+id & c-id \end{bmatrix} \begin{bmatrix} 1 & i \\ -1 & i \end{bmatrix} \\ &= \frac{1}{2i} \begin{bmatrix} 2ib & 2ia \\ 2id & 2ic \end{bmatrix} \\ &= \begin{bmatrix} b & a \\ d & c \end{bmatrix} \\ &= \begin{bmatrix} \frac{1}{\omega} & \frac{1}{v} \\ 1 & 1 \end{bmatrix} \end{aligned}$$

Thus we can find the change of basis matrix S w/o even knowing P or R . However, knowing P & remembering $R = \begin{bmatrix} i & -i \\ 1 & 1 \end{bmatrix}$ (always), we can verify our S .