


11.1: SEQUENCES

A is an ordered list  $a_1, a_2, \dots, a_n, \dots$

Common notations include

i)  $\{a_1, a_2, a_3, \dots\}$

ii)  $\{a_n\}$

iii)  $\{a_i\}_{i=1}^{\infty}$

w/ corresponding examples

i)  $\{1, 1, 2, 3, 5, \dots\}$

ii)  $\left\{\frac{(-1)^n}{n}\right\}$

iii)  $\left\{\frac{n!}{n^2}\right\}_{n=2}^{\infty}$

Ex 1: Find  $a_n$  if  $\{a_n\} = \left\{\frac{-2}{4}, \frac{3}{8}, \frac{-4}{16}, \dots\right\}$

What is the behavior of  $a_n = \frac{n}{n+1}$  as

$n \rightarrow \infty$ . Written another way,  $\lim_{n \rightarrow \infty} \frac{n}{n+1}$ .




What is  
the domain

Definition: A sequence  $\{a_n\}$  has the limit  $L$  if we can make the terms  $a_n$  arbitrarily close to  $L$  for sufficiently large  $n$ . Then we write  $\lim_{n \rightarrow \infty} a_n = L$ . Else,  $a_n$  diverges.

11.1
2/5

more precisely,

Definition. A sequence  $\{a_n\}$  converges to the limit  $L$  if  $\forall \epsilon > 0 \exists N$  s.t.  $a_n - L < \epsilon$  when  $n > N$ .



The only difference between this defn. and our old defn of the limit of a fct is the domain.

Theorem: If  $\lim_{x \rightarrow \infty} f(x) = L$  and  $f(n) = a_n$  when  $n \in \mathbb{Z}$ , then  $\lim_{n \rightarrow \infty} a_n = L$ .

Begin here day 2

Definition.  $\lim_{n \rightarrow \infty} a_n = \infty \Rightarrow \forall M > 0 \exists N \in \mathbb{Z}$  s.t.

$$n > N \Rightarrow a_n > M.$$

recall  $\forall \epsilon$

clarification  $\epsilon$  epsilon  
 $\in$  "is an element of."

The limit laws, if  $\{a_n\}$  and  $\{b_n\}$  are convergent sequences and  $c$  is a constant, then

11.1
3/5

$$i) \lim_{n \rightarrow \infty} (a_n \pm b_n) = \lim_{n \rightarrow \infty} a_n \pm \lim_{n \rightarrow \infty} b_n$$

$$ii) \lim_{n \rightarrow \infty} c a_n = c \lim_{n \rightarrow \infty} a_n$$

$$iii) \lim_{n \rightarrow \infty} (a_n b_n) = \lim_{n \rightarrow \infty} a_n \cdot \lim_{n \rightarrow \infty} b_n$$

$$iv) \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{\lim_{n \rightarrow \infty} a_n}{\lim_{n \rightarrow \infty} b_n} \quad \lim_{n \rightarrow \infty} b_n \neq 0$$

$$v) \lim_{n \rightarrow \infty} a_n^p = \left[ \lim_{n \rightarrow \infty} a_n \right]^p \quad \text{if } p > 0 \text{ and } a_n > 0$$

The Squeeze Theorem for sequences

If  $a_n \leq b_n \leq c_n$  for  $n \geq n_0$  and

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = L, \text{ then } \lim_{n \rightarrow \infty} b_n = L.$$

Theorem: If  $\lim_{n \rightarrow \infty} |a_n| = 0$ , then  $\lim_{n \rightarrow \infty} a_n = 0$

Ex2:  $\lim_{n \rightarrow \infty} \frac{\sin(n)}{n}$

l'Hopital's Rule on p 208.

EX3: Does  $\lim_{N \rightarrow \infty} \sin(N)$  converge or diverge.

11.1
4/5

EX4: Evaluate  $\lim_{N \rightarrow \infty} \frac{(-1)^N}{N^2}$  if it exists.

EX5: Evaluate  $\lim_{N \rightarrow \infty} \frac{N!}{N^N}$  if it exists.

$$\frac{N!}{N^N} = \frac{1}{N} \cdot \underbrace{\left( \frac{2 \cdot 3 \cdot \dots \cdot N}{N \cdot N \cdot \dots \cdot N} \right)}_{< 1}$$

so  $0 < \frac{N!}{N^N} < \frac{1}{N}$  AND the limit converges by the squeeze theorem.

ONLY IF time (hr).

EX6: Find the radius of convergence for  $\{n^n\}$

Defn.  $\{a_n\}$  is increasing if  $a_n < a_{n+1}$  for  $n \geq 1$  and decreasing if  $a_n > a_{n+1}$ .  $\{a_n\}$  is monotonic if it is increasing or decreasing.

EX7: show  $\{1/N\}$  is increasing.  
(use the derivative).

Defn.  $\{a_n\}$  is bounded <sup>above,</sup> if  $\exists M$  s.t.  $a_n \leq M \forall n \in \mathbb{N}$ . 11.1  
5/5

bounded below.  $a_n \geq m$ .

If  $\{a_n\}$  is bounded above and below,  
then we say it is a bounded sequence.

NOTE: upperbound vs. least upper bound.

Thm: Every bounded, monotonic sequence is  
convergent.

see proof in book ... completeness axiom.

NOTE: see CD for hints on real problems.

Hint: #65 & #72 are similar.