

## Complex Numbers.

$$z = a + bi$$

$|z|$  modulus.

$\theta$  argument of  $z$

$$\begin{aligned} z &= r \cos \theta + i r \sin \theta \\ &= r (\cos \theta + i \sin \theta). \end{aligned} \quad (\text{polar form of } z)$$

Thm: De Moivre's Thm

$$(\cos \theta + i \sin \theta)^n = \cos(n\theta) + i \sin(n\theta)$$

Rotations & Powers of  $z$ .

If  $z = r(\cos \theta + i \sin \theta)$ , then  $z^n$  spirals  
in/out as it rotates by  $\theta$  about the unit circle.

Thm: FTOA

Any poly  $p(\lambda)$  w/ complex coefficients splits,  
that is, it can be written as a product of  
linear factors.

$$p(\lambda) = k(\lambda - \lambda_1)(\lambda - \lambda_2) \dots (\lambda - \lambda_n)$$

for some complex numbers  $\lambda_1, \dots, \lambda_n$ , &  $k$ .

ex1: Diagonalize  $A = \begin{bmatrix} 11 & 6 \\ -15 & -7 \end{bmatrix}$

$$\begin{aligned} \det(A - \lambda I) &= \begin{vmatrix} 11 - \lambda & 6 \\ -15 & -7 - \lambda \end{vmatrix} \\ &= (11 - \lambda)(-7 - \lambda) + 90 \\ &= -77 - 4\lambda + \lambda^2 + 90 \\ &= \lambda^2 - 4\lambda + 13 \end{aligned}$$

Solve  $0 = \lambda^2 - 4\lambda + 13$

$$\begin{aligned} \lambda &= \frac{4 \pm \sqrt{16 - 4(1)(13)}}{2(1)} \\ &= \frac{4 \pm 6i}{2} \\ &= 2 \pm 3i \end{aligned}$$

Find  $\ker(A - (2 + 3i)I)$

$$\begin{bmatrix} 9 - 3i & 6 \\ -15 & -9 - 3i \end{bmatrix} \quad \frac{1}{9 - 3i} R_1 \rightarrow R_1$$

$$\begin{bmatrix} 1 & \frac{2}{3} + \frac{1}{3}i \\ -15 & -9 - 3i \end{bmatrix} \quad 15R_1 + R_2 \rightarrow R_2$$

$$\begin{bmatrix} 1 & \frac{2}{3} + \frac{1}{3}i \\ 0 & 0 \end{bmatrix}$$

$E_{2+3i} = \text{span} \left( \begin{bmatrix} -3 - i \\ 5 \end{bmatrix} \right)$

and check if the conjugate of  $\vec{v}_1$  is also an eigenvector.  $\lambda_2 \vec{v}_2$

$$\begin{bmatrix} 11 & 6 \\ -15 & -7 \end{bmatrix} \begin{bmatrix} -3 + i \\ 5 \end{bmatrix} = \begin{bmatrix} -3 + 11i \\ 10 - 15i \end{bmatrix}$$

So 
$$\begin{bmatrix} 2+3i & 0 \\ 0 & 2-3i \end{bmatrix} = P^{-1} A P \quad (\text{of the form } D = S^{-1} A S)$$

7.5  
3/4

where 
$$P = \begin{bmatrix} -3-i & -3+i \\ 5 & 5 \end{bmatrix}.$$

Again 
$$P^{-1} \begin{bmatrix} 11 & 6 \\ -15 & -7 \end{bmatrix} P = \begin{bmatrix} 2+3i & 0 \\ 0 & 2-3i \end{bmatrix}$$
 which is an example

reminding us that if there is an eigenbasis for the transformation  $T(\vec{x}) = A\vec{x}$  then  $A$  is diagonalizable.

In this case  $P$  or  $D$  contain complex entries.

ex2: Diagonalize  $A = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$

(rotation-scaling matrix)  
 $a, b \in \mathbb{R}$  and  $b \neq 0$ .

$\lambda = a \pm ib$

$E_{a+ib} = \text{span}$  It's very important that vectors are given as complex conjugates.

be careful — other eigenvectors don't work out. — see page 7.

and  $\mathbb{R}^2$

$\begin{bmatrix} 0 \\ a-ib \end{bmatrix}$

(of  $D$ ).

That is, we diagonalized scaling matrix.

recall: IF  $A$  is a real  $2 \times 2$  matrix w/ eigenvalues  $a \pm ib$  ( $b \neq 0$ ) and corresponding eigenvectors  $v \pm wi$ , then

$P^{-1} A P = \begin{bmatrix} a+ib & 0 \\ 0 & a-ib \end{bmatrix}$  where  $P = \begin{bmatrix} | & | \\ v+wi & v-wi \\ | & | \end{bmatrix}$

matrix w/ complex eigenvals.      rotation scaling matrix

$\Rightarrow P^{-1} A P = R^{-1} \begin{bmatrix} a & -b \\ b & a \end{bmatrix} R$

$\Rightarrow \begin{bmatrix} a & -b \\ b & a \end{bmatrix} = R P^{-1} A P R^{-1} = S^{-1} A S$

where  $S = P R^{-1} = \begin{bmatrix} | & | \\ v & w \\ | & | \end{bmatrix}$

$S$  has real values & so  $A$  is similar to a rotation-scaling matrix.

ex1 rev:  $A = \begin{bmatrix} 11 & 6 \\ -15 & -7 \end{bmatrix}$

$$\lambda = 2 \pm 3i$$

$$E_{2+3i} = \text{span} \left[ \begin{array}{c} -3 - i \\ 5 \end{array} \right]$$
  
$$\left[ \begin{array}{c} -3 \\ 5 \end{array} \right] + i \left[ \begin{array}{c} -1 \\ 0 \end{array} \right]$$
  
$$\vec{v} + i\vec{w}$$

$$\Rightarrow S = \begin{bmatrix} -1 & -3 \\ 0 & 5 \end{bmatrix}$$

and  $S^{-1} A S = \begin{bmatrix} 2 & -3 \\ 3 & 2 \end{bmatrix} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$

where  $\lambda = a + ib$  is an eigenvalue.

what scaling factor?

what rotation?

Thm: A complex  $n \times n$  matrix has  $n$  complex eigenvals if they are counted w/ alg. mult.

Thm:  $\det(A) = \lambda_1 \dots \lambda_n$

$$\text{Tr}(A) = \lambda_1 + \dots + \lambda_n$$

## THE FORMULA FOR S

7.5  
6/7

$$\text{If } R = \begin{bmatrix} i & -i \\ 1 & 1 \end{bmatrix} \text{ and } P = \begin{bmatrix} \bar{v} + i\bar{w} & v - i\bar{w} \\ 1 & 1 \end{bmatrix}$$

$$\begin{aligned} \text{Then } S = PR^{-1} &= \frac{1}{2i} \begin{bmatrix} \bar{v} + i\bar{w} & v - i\bar{w} \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & i \\ -1 & i \end{bmatrix} \\ &= \frac{1}{2i} \begin{bmatrix} a + ib & a - ib \\ c + id & c - id \end{bmatrix} \begin{bmatrix} 1 & i \\ -1 & i \end{bmatrix} \\ &= \frac{1}{2i} \begin{bmatrix} 2ib & 2ia \\ 2id & 2ic \end{bmatrix} \\ &= \begin{bmatrix} b & a \\ d & c \end{bmatrix} \\ &= \begin{bmatrix} \frac{1}{\bar{w}} & \frac{1}{\bar{v}} \\ 1 & 1 \end{bmatrix} \end{aligned}$$

Thus we can find the change of basis matrix  $S$  w/o even knowing  $P$  or  $R$ . However, knowing  $P$  & remembering  $R = \begin{bmatrix} i & -i \\ 1 & 1 \end{bmatrix}$  (always), we can verify our  $S$ .

# CHANGE OF BASIS MATRICES REQUIRE COMPLEX CONJUGATE EIGENVECTOR

$\lambda = 2 \pm 3i$  for the matrices  $A = \begin{bmatrix} 11 & 6 \\ -15 & -7 \end{bmatrix}$  and  $P = \begin{bmatrix} -3-i & -3+i \\ 5 & 5 \end{bmatrix}$

Eigenvectors of  $B = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$  can take many forms. 7.5  
7/7

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$\vec{v}_2 = \begin{bmatrix} -i \\ 1 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} i \\ -1 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} 1 \\ i \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} -2i \\ 2 \end{bmatrix}$$

$\vec{v}_1$  conjugate      \*(-1)      \*(i)      \*(2)

which gives R many forms

$$R = \begin{bmatrix} i & -i \\ 1 & 1 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} i & i \\ 1 & -1 \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} i & 1 \\ 1 & i \end{bmatrix} \quad \text{or} \quad \begin{bmatrix} i & -2i \\ 1 & 2 \end{bmatrix}$$

check R & P by comparing the product

$$S = PR^{-1} \quad \text{to the formula for } S = \begin{bmatrix} \vec{u} & \vec{v} \end{bmatrix}$$

where the eigenvectors of  $A = \vec{v} + i\vec{w} = \begin{bmatrix} -3 \\ 5 \end{bmatrix} \pm i \begin{bmatrix} 4 \\ 0 \end{bmatrix}$

$$\vec{v}_1 \text{ conjugate: } PR^{-1} = \frac{1}{2i} \begin{bmatrix} -3-i & -3+i \\ 5 & 5 \end{bmatrix} \begin{bmatrix} 1 & i \\ -1 & i \end{bmatrix} = \frac{1}{2i} \begin{bmatrix} -2i & -6i \\ 0 & 10i \end{bmatrix} = \begin{bmatrix} -1 & -3 \\ 0 & 5 \end{bmatrix} = S$$

$$*(-1): PR^{-1} = -\frac{1}{2i} \begin{bmatrix} -3-i & -3+i \\ 5 & 5 \end{bmatrix} \begin{bmatrix} -i & -i \\ -1 & i \end{bmatrix} = +\frac{1}{2i} \begin{bmatrix} -6 & +2 \\ +10 & 0 \end{bmatrix} = \begin{bmatrix} 3i & -i \\ -5i & 0 \end{bmatrix} \neq S$$

$$*(i): PR^{-1} = -\frac{1}{2} \begin{bmatrix} -3-i & -3+i \\ 5 & 5 \end{bmatrix} \begin{bmatrix} i & -1 \\ -1 & i \end{bmatrix} = -\frac{1}{2} \begin{bmatrix} -4i+4 & 2-2i \\ 5i-5 & -5+5i \end{bmatrix} \neq S$$

$$*(2): PR^{-1} = \frac{1}{4i} \begin{bmatrix} -3-i & -3+i \\ 5 & 5 \end{bmatrix} \begin{bmatrix} 2 & 2i \\ -1 & i \end{bmatrix} = \frac{1}{4i} \begin{bmatrix} -3-3i & -9i+1 \\ 5 & 15i \end{bmatrix} \neq S$$

Ex:

$$A = \begin{bmatrix} 4 & 2 \\ -1 & 2 \end{bmatrix}$$

eigenvalues:  $\lambda = 3+i$  and  $\lambda = 3-i$

eigenvectors:  $\vec{v}_1 = \begin{bmatrix} -1-i \\ 1 \end{bmatrix}$  and  $\vec{v}_2 = \begin{bmatrix} -1+i \\ 1 \end{bmatrix}$

$$\begin{array}{c} \uparrow \\ \begin{bmatrix} -1 \\ 1 \end{bmatrix} + i \begin{bmatrix} -1 \\ 0 \end{bmatrix} \\ \vec{v} + i\vec{w} \end{array}$$

$$A = S B S^{-1} \quad \text{w/} \quad S = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ i & -i \end{bmatrix} = \begin{bmatrix} -1 & -1 \\ \Delta & 1 \end{bmatrix}$$

$$B = \begin{bmatrix} 3 & -1 \\ 1 & 3 \end{bmatrix}$$