Concept Check

- 1. What is the difference between a vector and a scalar?
- 2. How do you add two vectors geometrically? How do you add them algebraically?
- **3.** If **a** is a vector and *c* is a scalar, how is *c***a** related to **a** geometrically? How do you find *c***a** algebraically?
- 4. How do you find the vector from one point to another?
- 5. How do you find the dot product $\mathbf{a} \cdot \mathbf{b}$ of two vectors if you know their lengths and the angle between them? What if you know their components?
- 6. How are dot products useful?
- Write expressions for the scalar and vector projections of b onto a. Illustrate with diagrams.
- 8. How do you find the cross product $\mathbf{a} \times \mathbf{b}$ of two vectors if you know their lengths and the angle between them? What if you know their components?
- 9. How are cross products useful?
- 10. (a) How do you find the area of the parallelogram determined by a and b?
 - (b) How do you find the volume of the parallelepiped determined by **a**, **b**, and **c**?

- 11. How do you find a vector perpendicular to a plane?
- 12. How do you find the angle between two intersecting planes?
- **13.** Write a vector equation, parametric equations, and symmetric equations for a line.
- 14. Write a vector equation and a scalar equation for a plane.
- 15. (a) How do you tell if two vectors are parallel?
 - (b) How do you tell if two vectors are perpendicular?
 - (c) How do you tell if two planes are parallel?
- **16.** (a) Describe a method for determining whether three points P, Q, and R lie on the same line.
 - (b) Describe a method for determining whether four points P, Q, R, and S lie in the same plane.
- 17. (a) How do you find the distance from a point to a line?
 - (b) How do you find the distance from a point to a plane?
 - (c) How do you find the distance between two lines?
- 18. What are the traces of a surface? How do you find them?
- **19.** Write equations in standard form of the six types of quadric surfaces.

True-False Quiz

Determine whether the statement is true or false. If it is true, explain why. If it is false, explain why or give an example that disproves the statement.

- 1. If $\mathbf{u} = \langle u_1, u_2 \rangle$ and $\mathbf{v} = \langle v_1, v_2 \rangle$, then $\mathbf{u} \cdot \mathbf{v} = \langle u_1 v_1, u_2 v_2 \rangle$.
- **2.** For any vectors **u** and **v** in V_3 , $|\mathbf{u} + \mathbf{v}| = |\mathbf{u}| + |\mathbf{v}|$.
- **3.** For any vectors \mathbf{u} and \mathbf{v} in V_3 , $|\mathbf{u} \cdot \mathbf{v}| = |\mathbf{u}| |\mathbf{v}|$.
- **4.** For any vectors \mathbf{u} and \mathbf{v} in V_3 , $|\mathbf{u} \times \mathbf{v}| = |\mathbf{u}| |\mathbf{v}|$.
- 5. For any vectors \mathbf{u} and \mathbf{v} in V_3 , $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$.

- 6. For any vectors \mathbf{u} and \mathbf{v} in V_3 , $\mathbf{u} \times \mathbf{v} = \mathbf{v} \times \mathbf{u}$.
- 7. For any vectors \mathbf{u} and \mathbf{v} in V_3 , $|\mathbf{u} \times \mathbf{v}| = |\mathbf{v} \times \mathbf{u}|$.
- 8. For any vectors \mathbf{u} and \mathbf{v} in V_3 and any scalar k, $k(\mathbf{u} \cdot \mathbf{v}) = (k\mathbf{u}) \cdot \mathbf{v}$.
- **9.** For any vectors \mathbf{u} and \mathbf{v} in V_3 and any scalar k, $k(\mathbf{u} \times \mathbf{v}) = (k\mathbf{u}) \times \mathbf{v}$.
- **10.** For any vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in V_3 , $(\mathbf{u} + \mathbf{v}) \times \mathbf{w} = \mathbf{u} \times \mathbf{w} + \mathbf{v} \times \mathbf{w}$.

859

- 11. For any vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in V_3 , $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = (\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}$
- 12. For any vectors \mathbf{u} , \mathbf{v} , and \mathbf{w} in V_3 , $\mathbf{u} \times (\mathbf{v} \times \mathbf{w}) = (\mathbf{u} \times \mathbf{v}) \times \mathbf{w}$.
- 13. For any vectors \mathbf{u} and \mathbf{v} in V_3 , $(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{u} = 0$.
- 14. For any vectors \mathbf{u} and \mathbf{v} in V_3 , $(\mathbf{u} + \mathbf{v}) \times \mathbf{v} = \mathbf{u} \times \mathbf{v}$.
- **15.** The vector $\langle 3, -1, 2 \rangle$ is parallel to the plane 6x - 2y + 4z = 1.

- **16.** A linear equation Ax + By + Cz + D = 0 represents a line in space,
- 17. The set of points $\{(x, y, z) \mid x^2 + y^2 = 1\}$ is a circle.
- **18.** In \mathbb{R}^3 the graph of $y = x^2$ is a paraboloid.
- 19. If $\mathbf{u} \cdot \mathbf{v} = 0$, then $\mathbf{u} = \mathbf{0}$ or $\mathbf{v} = \mathbf{0}$.
- 20. If $\mathbf{u} \times \mathbf{v} = \mathbf{0}$, then $\mathbf{u} = \mathbf{0}$ or $\mathbf{v} = \mathbf{0}$.
- 21. If $\mathbf{u} \cdot \mathbf{v} = 0$ and $\mathbf{u} \times \mathbf{v} = \mathbf{0}$, then $\mathbf{u} = \mathbf{0}$ or $\mathbf{v} = \mathbf{0}$.
- 22. If \mathbf{u} and \mathbf{v} are in V_3 , then $|\mathbf{u} \cdot \mathbf{v}| \leq |\mathbf{u}| |\mathbf{v}|$.


Exercises

- 1. (a) Find an equation of the sphere that passes through the point (6, -2, 3) and has center (-1, 2, 1).
 - (b) Find the curve in which this sphere intersects the yz-plane.
 - (c) Find the center and radius of the sphere


$$x^2 + y^2 + z^2 - 8x + 2y + 6z + 1 = 0$$

- 2. Copy the vectors in the figure and use them to draw each of the following vectors.
 - (a) $\mathbf{a} + \mathbf{b}$
- (b) $\mathbf{a} \mathbf{b}$

- (d) $2\mathbf{a} + \mathbf{b}$



3. If u and v are the vectors shown in the figure, find $u\,\cdot\,v$ and $|\mathbf{u} \times \mathbf{v}|$. Is $\mathbf{u} \times \mathbf{v}$ directed into the page or out of it?



4. Calculate the given quantity if

$$\mathbf{a} = \mathbf{i} + \mathbf{i} - 2\mathbf{k}$$

$$\mathbf{b} = 3\mathbf{i} - 2\mathbf{j} + \mathbf{k}$$

$$c = j - 5k$$


- (a) 2a + 3b
- (b) **b**
- (c) a · b
- (d) $\mathbf{a} \times \mathbf{b}$
- (e) $|\mathbf{b} \times \mathbf{c}|$
- (f) $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$
- $(g) c \times c$
- (b) $\mathbf{a} \times (\mathbf{b} \times \mathbf{c})$
- (i) compa b
- (j) proja b

- (k) The angle between a and b (correct to the nearest degree)
- 5. Find the values of x such that the vectors (3, 2, x) and $\langle 2x, 4, x \rangle$ are orthogonal.


- **6.** Find two unit vectors that are orthogonal to both $\mathbf{j} + 2\mathbf{k}$ and i - 2j + 3k.
- 7. Suppose that $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = 2$. Find
 - (a) $(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}$
- (b) $\mathbf{u} \cdot (\mathbf{w} \times \mathbf{v})$
- (c) $\mathbf{v} \cdot (\mathbf{u} \times \mathbf{w})$
- (d) $(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{v}$
- **8.** Show that if \mathbf{a} , \mathbf{b} , and \mathbf{c} are in V_3 , then

$$(\mathbf{a} \times \mathbf{b}) \cdot [(\mathbf{b} \times \mathbf{c}) \times (\mathbf{c} \times \mathbf{a})] = [\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})]^2$$

- 9. Find the acute angle between two diagonals of a cube.
- **10.** Given the points A(1, 0, 1), B(2, 3, 0), C(-1, 1, 4), and D(0, 3, 2), find the volume of the parallelepiped with adjacent edges AB, AC, and AD.
- 11. (a) Find a vector perpendicular to the plane through the points A(1, 0, 0), B(2, 0, -1), and C(1, 4, 3).
 - (b) Find the area of triangle ABC.
- 12. A constant force $\mathbf{F} = 3\mathbf{i} + 5\mathbf{j} + 10\mathbf{k}$ moves an object along the line segment from (1, 0, 2) to (5, 3, 8). Find the work done if the distance is measured in meters and the force in newtons.
- 13. A boat is pulled onto shore using two ropes, as shown in the diagram. If a force of 255 N is needed, find the magnitude of the force in each rope.



14. Find the magnitude of the torque about P if a 50-N force is applied as shown.



- **15.** The line through (4, -1, 2) and (1, 1, 5)
- **16.** The line through (1, 0, -1) and parallel to the line $\frac{1}{3}(x 4) = \frac{1}{2}y = z + 2$
- 17. The line through (-2, 2, 4) and perpendicular to the plane 2x y + 5z = 12
- 18-20 Find an equation of the plane.
- **18.** The plane through (2, 1, 0) and parallel to x + 4y 3z = 1
- **19.** The plane through (3, -1, 1), (4, 0, 2), and (6, 3, 1)
- **20.** The plane through (1, 2, -2) that contains the line x = 2t, y = 3 t, z = 1 + 3t
- 21. Find the point in which the line with parametric equations x = 2 t, y = 1 + 3t, z = 4t intersects the plane 2x y + z = 2.
- 22. Find the distance from the origin to the line x = 1 + t, y = 2 t, z = -1 + 2t.
- **23.** Determine whether the lines given by the symmetric equations

$$\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4}$$

and

$$\frac{x+1}{6} = \frac{y-3}{-1} = \frac{z+5}{2}$$

are parallel, skew, or intersecting.

24. (a) Show that the planes x + y - z = 1 and 2x - 3y + 4z = 5 are neither parallel nor perpendicular.

- (b) Find, correct to the nearest degree, the angle between these planes.
- 25. Find an equation of the plane through the line of intersection of the planes x z = 1 and y + 2z = 3 and perpendicular to the plane x + y 2z = 1.
- **26.** (a) Find an equation of the plane that passes through the points A(2, 1, 1), B(-1, -1, 10), and C(1, 3, -4).
 - (b) Find symmetric equations for the line through B that is perpendicular to the plane in part (a).
 - (c) A second plane passes through (2, 0, 4) and has normal vector (2, -4, -3). Show that the acute angle between the planes is approximately 43° .
 - (d) Find parametric equations for the line of intersection of the two planes.
- 27. Find the distance between the planes 3x + y 4z = 2 and 3x + y 4z = 24.
- 28-36 Identify and sketch the graph of each surface.

28.
$$x = 3$$

29.
$$x = z$$

30.
$$v = z^2$$

31.
$$x^2 = y^2 + 4z^2$$

32.
$$4x - y + 2z = 4$$

33.
$$-4x^2 + y^2 - 4z^2 = 4$$

34.
$$v^2 + z^2 = 1 + x^2$$

35.
$$4x^2 + 4y^2 - 8y + z^2 = 0$$

36.
$$x = y^2 + z^2 - 2y - 4z + 5$$

- **37.** An ellipsoid is created by rotating the ellipse $4x^2 + y^2 = 16$ about the *x*-axis. Find an equation of the ellipsoid.
- **38.** A surface consists of all points P such that the distance from P to the plane y = 1 is twice the distance from P to the point (0, -1, 0). Find an equation for this surface and identify it.

CHAPTER 12 REVIEW # PAGE 858

True-False Quiz


- 9. True 7. True 5. True 1. False 3. False
- 15. False 17. False 13. True 11. True
- **21**. True 19. False


Exercises

- 1. (a) $(x + 1)^2 + (y 2)^2 + (z 1)^2 = 69$
- (b) $(y-2)^2 + (z-1)^2 = 68, x = 0$


- (c) Center (4, -1, -3), radius 5 3. $\mathbf{u} \cdot \mathbf{v} = 3\sqrt{2}$; $|\mathbf{u} \times \mathbf{v}| = 3\sqrt{2}$; out of the page 5. -2, -4 7. (a) 2 (b) -2 (c) -2 (d) 0
- **9.** $\cos^{-1}(\frac{1}{3}) \approx 71^{\circ}$ **11.** (a) $\langle 4, -3, 4 \rangle$ (b) $\sqrt{41/2}$
- **13.** 166 N, 114 N
- 15. x = 4 3t, y = -1 + 2t, z = 2 + 3t
- 17. x = -2 + 2t, y = 2 t, z = 4 + 5t
- **19.** -4x + 3y + z = -14 **21.** (1, 4, 4) 23. Skew
- **25.** x + y + z = 4 **27.** $22/\sqrt{26}$
- **29**. Plane


31. Cone





33. Hyperboloid of two sheets 35. Ellipsoid





37. $4x^2 + y^2 + z^2 = 16$

PROBLEMS PLUS = PAGE 861

- 1. $(\sqrt{3} \frac{3}{2})$ m
- 3. (a) $(x+1)/(-2c) = (y-c)/(c^2-1) = (z-c)/(c^2+1)$ (b) $x^2 + y^2 = t^2 + 1$, z = t (c) $4\pi/3$
- **5**: 20