

$$\int \sqrt{a^2 - x^2} dx$$

$$\text{Let } x = a \sin(\theta) \quad \text{on } -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$$

$$= \sqrt{a^2 - a^2 \sin^2(\theta)}$$

$$= \sqrt{a^2 (1 - \sin^2(\theta))}$$

$$= \sqrt{a^2 \cos^2(\theta)}$$

$$= a |\cos(\theta)| \quad \text{but } \cos(\theta) \geq 0 \quad \text{on } -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$$

$$= a \cos(\theta).$$

$$\text{Ex 1: } \int x^2 \sqrt{4-x^2} dx$$

$$\text{Let } x = 2 \sin(\theta) \Rightarrow dx = 2 \cos(\theta) d\theta.$$

$$= \int 4 \sin^2(\theta) \sqrt{4 - 4 \sin^2(\theta)} \cdot 2 \cos(\theta) d\theta$$

$$= \int 4 \sin^2(\theta) (2 \cos(\theta)) \cdot 2 \cos(\theta) d\theta$$

$$= 16 \int \sin^2(\theta) \cdot \cos^2(\theta) d\theta.$$


$$= 16 \int \left( \frac{1 - \cos(2\theta)}{2} \right) \left( \frac{1 + \cos(2\theta)}{2} \right) d\theta$$

$$= 4 \int [1 - \cos^2(2\theta)] d\theta$$

2

$$\begin{aligned}
 \hookrightarrow &= 4 \int \left[ 1 - \frac{1 + \cos(4\theta)}{2} \right] d\theta \\
 &= 2 \int [2 - 1 - \cos(4\theta)] d\theta \\
 &= 2 \int [1 - \cos(4\theta)] d\theta \\
 &= 2 \left[ \theta - \frac{\sin(4\theta)}{4} \right] + C \\
 &= 2 \left[ \theta - \frac{1}{4} (2 \sin(2\theta) \cos(2\theta)) \right] + C \\
 &= 2 \left[ \theta - \frac{1}{4} (4 \sin(\theta) \cos(\theta) (\cos^2(\theta) - \sin^2(\theta))) \right] + C
 \end{aligned}$$

since  $x = 2 \sin(\theta) \Rightarrow \frac{x}{2} = \sin(\theta)$ .



$$= 2 \left[ \sin^{-1}\left(\frac{x}{2}\right) + \frac{x}{2} \left(\frac{\sqrt{4-x^2}}{2}\right) \left(\frac{4-x^2}{4} - \frac{x^2}{4}\right) \right] + C$$

⋮

To work w/ expressions...

expression	Sub	Identity
$\sqrt{a^2 - x^2}$	$x = a \sin(\theta), \quad -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$	$1 - \sin^2(\theta) = \cos^2$
$\sqrt{a^2 + x^2}$	$x = a \tan(\theta)$	$1 + \tan^2 = \sec^2$
$\sqrt{x^2 - a^2}$	$x = a \sec(\theta), \quad \theta \leq \theta < \frac{\pi}{2} \text{ or } \pi \leq \theta < \frac{3\pi}{2}$	$\sec^2 - 1 = \tan^2$

Ex 2: Find the area of a circle w/ radius a.

Ex 3:  ~~$\int \dots$~~   $\int x \sqrt{x^2 + 16} dx$

Ex 4:  $\int \frac{1}{\sqrt{9x^2 + 6x - 8}} dx$  complete the square.

Ex 5:  $\int \frac{dx}{(x^2 - 1)^{3/2}} = \int \frac{dx}{(\sqrt{x^2 - 1})^3}$  Let  $x = \sec(\theta)$ .  
 $dx = \sec(\theta) \tan(\theta) d\theta$   
 $= \int \frac{\sec(\theta) \tan(\theta)}{(\sqrt{\sec^2(\theta) - 1})^3} d\theta$   
 $= \int \frac{\sec(\theta) \tan(\theta)}{\tan^3(\theta)} d\theta$   
 $= \int \frac{\sec(\theta)}{\tan^2(\theta)} d\theta = \int \frac{\cos^2(\theta)}{\sin^2(\theta) \cos(\theta)} d\theta$