Math 220: Linear Algebra

Linear Equations

A linear of the variables $x_1, x_2, ..., x_n$ has the form $a_1x_1 + a_2x_2 + ... + a_nx_n = b$

Where $b, a_1, a_2, ..., a_n$ are real or complex numbers.

Ex 1: Circle the linear equations, and state why the non-linear aren't linear.

(a)
$$3x_1 - x_2 = 5x_1$$
 b) $4x_2 + 5 = \sqrt{x_1}$ b) $4x_2 + 5 = \sqrt{x_1}$ b) $4x_2 + 5 = \sqrt{x_1}$ c) $x_1 - 4x_2 = x_1x_2$ d) $-2x_1 + 7x_2 - \pi x_3 = \sqrt{2}$

A <u>System</u> of <u>livear</u> <u>equations</u> is a collection of one or more linear equations with the same variables. For example

$$3x_1 - x_2 - 4x_3 = 3$$
$$x_1 - 5x_3 = -2$$


A ______ of a system is a list of numbers $(s_1, s_2, ..., s_n)$ that make every equation of the system true, when each s_k is substituted for x_k .

The set of all possible solutions is called the <u>Solution</u> <u>set</u>.

Ex 3: Find another solution to the system from Ex 2.

Two systems are considered <u>equivalent</u> if they have the same solution set.

From 2 dimensional systems of equations in algebra, we should remember that there are 3 possibilities for the number of solutions to a system.



A system of linear equations has

- 1. no solution, or
- 2. exactly one solution, or
- 3. infinitely many solutions.

A system is called _______ if it has at least one solution, and ______ if it has no solutions.

Matrix Notation

We will represent a system of equations by its coefficients in a _____ へるナゲラス

$$x_1 -3x_3 = 8 \qquad \text{will be re-written as the} \\ 2x_1 + 2x_2 + 9x_3 = 7 \\ x_2 + 5x_3 = -2$$

The <u>size</u> of a matrix tells how many <u>rows</u> and <u>colorings</u> a matrix has.

An $m \times n$ matrix has

m rows and works.

Page 2 of 4

Solving a Linear System – We are going to describe an algorithm for solving linear systems, which replaces one system with an equivalent one that is easier to solve. Since they are equivalent, they have the same solution set.

Ex 4: Solve the system

Elementary Row Operations

- 1. (Replacement) Replace one row by the sum of itself and a multiple of another row.
- 2. (Interchange) Interchange two rows.
- 3. (Scaling) Multiply all entries in a row by a nonzero constant.

Two matrices are called <u>row</u> <u>equivalent</u> if there are a sequence of elementary row operations that transform one matrix into the other.

If two systems are row equivalent, they have the same ______ <u> らのいれのい </u> <u> とみ、</u>.

Two Fundamental Questions About a Linear System

- 1. Is the system consistent; that is, does at least one solution exist?
- 2. If a solution exists, is it the only one; that is, is the solution unique?

Ex 5: Determine whether the systems are consistent or inconsistent. Do not solve.

$$x_{2} + 4x_{3} = -5$$

$$x_{1} + 3x_{2} + 5x_{3} = -2$$

$$3x_{1} + 7x_{2} + 7x_{3} = 6$$

$$\begin{bmatrix} 0 & 1 & 4 & -5 \\ 1 & 5 & 5 & -2 \\ 5 & 7 & 7 & 6 \end{bmatrix} R_{1} + R_{2}$$

$$\begin{bmatrix} 1 & 3 & 5 & -2 \\ 0 & 1 & 4 & -5 \\ 3 & 7 & 7 & 6 \end{bmatrix} R_{3} - 3R_{1} - 3R_{3}$$

$$\begin{bmatrix} 1 & 3 & 5 & -2 \\ 0 & 1 & 4 & -5 \\ 0 & -2 & -8 & 12 \end{bmatrix} R_{3} + 2R_{2} - R_{3}$$

$$\begin{bmatrix} 1 & 3 & 5 & -2 \\ 0 & 1 & 4 & -5 \\ 0 & -2 & -8 & 12 \end{bmatrix} R_{3} + 2R_{2} - R_{3}$$

$$\begin{bmatrix} 1 & 3 & 5 & -2 \\ 0 & 1 & 4 & -5 \\ 0 & 0 & 2 \end{bmatrix}$$
This means $0 = 2$
The system is Page 4 of 4

$$x_{1} + 3x_{3} = 2$$

$$x_{2} - 3x_{4} = 3$$

$$-2x_{2} + 3x_{3} + 2x_{4} = 1$$

$$3x_{1} + 7x_{4} = -5$$

$$\begin{bmatrix} 1 & 0 & 3 & 0 & 2 \\ 0 & 2 & 0 & -3 & 3 \\ 0 & -2 & 3 & 2 & 1 \\ 2 & 0 & 0 & 7 & -5 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 3 & 0 & 2 \\ 0 & 2 & 0 & -3 & 3 \\ 0 & -2 & 3 & 2 & 1 \\ 0 & 0 & -4 & 7 & -11 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 3 & 0 & 2 \\ 0 & 2 & 0 & -3 & 3 \\ 0 & 0 & -4 & 7 & -11 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 3 & 0 & 2 \\ 0 & 2 & 0 & -3 & 3 \\ 0 & 0 & 7 & -11 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 3 & 0 & 2 \\ 0 & 2 & 0 & -3 & 3 \\ 0 & 0 & 7 & -11 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 3 & 0 & 2 \\ 0 & 2 & 0 & -3 & 3 \\ 0 & 0 & 7 & -11 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 3 & 0 & 2 \\ 0 & 2 & 0 & -3 & 3 \\ 0 & 0 & 7 & -11 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 3 & 0 & 2 \\ 0 & 2 & 0 & -3 & 3 \\ 0 & 0 & 7 & -11 \end{bmatrix}$$
There will be values for $X_{1,1,1,1}, X_{2}$